

Elsevier Editorial System(tm) for CORTEX  
Manuscript Draft

Manuscript Number: CORTEX-D-09-00028R1

Title: The mechanism of Transcranial Magnetic Stimulation in cognition

Article Type: Special Issue - Rusconi and Bestmann

Keywords: Transcranial magnetic stimulation; rTMS; noise; virtual lesion; cognition

Corresponding Author: Dr Carlo Miniussi,

Corresponding Author's Institution:

First Author: Carlo Miniussi

Order of Authors: Carlo Miniussi; Manuela Ruzzoli; Vincent Walsh

Abstract: As this is an opinion paper, no abstract is provided

Response to Reviewers: see file

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## The mechanism of Transcranial Magnetic Stimulation in cognition

Carlo Miniussi<sup>1,2</sup>, Manuela Ruzzoli<sup>2,3</sup>, Vincent Walsh<sup>4</sup>

<sup>1</sup>Department of Biomedical Sciences and Biotechnology, National Institute of Neuroscience, University of Brescia, Italy.

<sup>2</sup>Cognitive Neuroscience Section, IRCCS San Giovanni di Dio Fatebenefratelli, Brescia, Italy.

<sup>3</sup>Department of Neurological and Vision Sciences, University of Verona, Verona, Italy.

<sup>4</sup>Institute of Cognitive Neuroscience and Department of Psychology, University College London, London UK.

Corresponding author: Carlo Miniussi

Tel. +39 0303717441; fax. +39 0303717443; e-mail: [miniussi@med.unibs.it](mailto:miniussi@med.unibs.it)

Despite the widespread usage of transcranial magnetic stimulation (TMS) in clinical and basic research, the exact mechanisms of action and interactions with ongoing neural activity remain unclear. However thanks to recent biophysical studies on electromagnetic induction of neural tissue (Wagner et al., in press) we now know more about some basic properties of TMS effects. This basic knowledge is important in planning and interpreting TMS studies and in cognitive neuroscience experiments a theoretical framework is also necessary.

TMS data have traditionally been interpreted in the “virtual brain lesion” framework (Walsh and Cowey 1998). The terminology was proposed by analogy with neuropsychological and animal lesion studies and TMS is described as inducing a temporary, reversible lesion in the stimulated area, avoiding problems related to cortical plasticity and functional reorganisation (Walsh and Pascual-Leone 2003). Based on this interpretation, TMS has been utilised to define the putative role of areas during the execution of cognitive tasks, and this approach has been very productive.

1 Semantically, the term “virtual lesion” identifies the effect induced by TMS as blocking the  
2 function of a population of neurons that are temporarily “lesioned” by the TMS pulse. Nevertheless,  
3 the TMS pulse induces a depolarisation of a group of neurons that in turn might activate other  
4 neurons, and the final behavioural outcome depends on the role of the stimulated area in relation to  
5 the network engaged in such a task (Sack and Linden 2003). The “virtual lesion” hypothesis has,  
6 however, confused people on the issue of how TMS can possibly lead to enhanced performance  
7 (e.g., Harris et al., 2008b; Walsh et al., 1998). In addition, the brain may also compensate for  
8 interference either within an area or across a circuit because it does not react passively to cortical  
9 stimulation and because the state of activation and the task demand influence the response (e.g.,  
10 Bestmann et al., 2008; Ruff et al., in press; Siebner et al., in press; Silvanto et al., 2008). Finally, the  
11 virtual lesion term is just that, words, and it is not informative about the possible mechanisms of  
12 action of TMS. It is unclear whether one best describes TMS as suppressing neural signals, or if it  
13 adds random neural activity in the stimulated area (Walsh and Cowey 2000; Harris et al., 2008a).  
14 The result in both cases will be altered information processing, but we have one “hypothesis” (if for  
15 the sake of argument we can aggrandise the analogy in that way) and (at least) two possible  
16 mechanisms of action (i.e., either suppression of the relevant signal or addition of random neural  
17 noise).

18 For these reasons, we now have a choice: a separate explanation for each disruption or enhancement,  
19 or a mechanistic explanation of one effect producing positive or negative effects depending on the  
20 task, timing of TMS and areas involved. Nevertheless, it seems that because a great deal of TMS  
21 research does adopt a “point and shoot” methodology and conceptualises results as negative  
22 (“lesion”) or positive (“paradoxical”) it is clear that some change in terminology or  
23 reconceptualization of the effects of TMS should be discussed. Interpretations of data can go  
24 beyond a simple relationship between an anatomical area and impairment of behaviour as suggested  
25 by the virtual lesion terminology.

1 The key point is that we need to dissociate the language of physiological effects from those of  
2 behavioural effects (Miniussi et al., 2008; some papers unfortunately term disruptions as “due to  
3 inhibition” and facilitations as “due to excitation”). Our suggestion here is that behavioural effects  
4 are described in terms of models of the psychological measures. The best available model of  
5 decisions for most psychological experiments is signal detection theory (SDT: Green and Swets  
6 1966) and this allows one to dissociate statements of effects from premature statements about  
7 mechanism. The behavioural outcome of a system (e.g., reaction times or accuracy) represents the  
8 final output of a population of neurons whose activity is based on the functional attitude of the  
9 activated system (i.e., its morphology and state) and the task demands (i.e., input). Those neurons  
10 that respond in the same way to the task-goal will display signal correlation, represented as similar  
11 tuning curves that will contribute to the final output (Stein et al., 2005). There is also another source  
12 of activity that does not contribute to the goal and will determine the trial-to-trial response  
13 variability (Stein et al., 2005). We can define this activity as *neuronal noise*, a term that describes  
14 the “random” activity of neurons that is not associated with the encoding of behaviourally relevant  
15 variables. Noise is generally considered the main factor that limits the capacity of information  
16 processing by the brain and its effects could be of theoretical interest in a biological context (e.g.,  
17 Ermentrout et al., 2008; Stein et al., 2005). In general, noise decreases performance, but nonlinear  
18 systems, like the brain, can use noise to enhance performance through stochastic resonance (Moss et  
19 al., 2004). The presence of neuronal noise might confer on neurons more sensitivity to a given  
20 range of weak inputs, i.e., those neurons “randomly activated” and that go in the same direction of  
21 the signal, thereby rendering the noise in the signal (Stein et al., 2005). Noise can either interfere  
22 with or facilitate performance. It is easy to imagine how random noise can interfere with  
23 performance, for example by either introducing temporally random firing or obstructing  
24 synchronised interactions between populations. The opposite may be observed if the induced  
25 activity is somehow synchronised with the temporal coding between neural populations (i.e., neural

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

synchronisation or phase locking) at an adequate intensity (i.e., optimum level) (Ermentrout et al., 2008; Stein et al., 2005).

We can now return to how to interpret TMS effects produced by interactions with ongoing neural activity. We could refer to this activity as neural noise because it is activity artificially induced in the system. Thus, TMS could modify system activity by altering whatever information is carried by a precise induced neural firing pattern. This effect could be interpreted as the TMS generating neural activity (noise) with respect to the relevant information carried by the stimulated area (signal). Nevertheless, TMS will influence not only neurons that contribute to noise but also those that code for correct responses (i.e., the finalised activity, or signal). TMS may induce neuronal activity that adds to the *ongoing neural activity* as a complement to the extant activity pattern (i.e., activated by state and task demand), which can be considered both as noise and as part of the signal, depending on the neuron population that will be activated.

In this framework, it is also possible to explain facilitatory results in terms of the strict relationship between noise and signal in the nervous system: enhanced performance, for example, may be observed with an optimum level of noise. It is also possible to explain different effects induced in the same area, depending on the task demands and stimulation parameters. Thus, the relationship between signal and noise seems to offer a better view of the induced effects when considering that both have the same nature (neural activity) and that they provide complementary rather than mutually exclusive information.

It is clearly important that we pursue more precise descriptions of TMS-induced effects to allow more sophisticated inferences (beyond area X is necessary for function Y). Such a theoretical framework is necessary to interpret the effects of double dissociations, cortico-cortical interaction, state and plasticity and will therefore improve the range of parameters at our disposal in programmes of rehabilitation. Such an opportunity to have a formal framework in which precise

1 behavioural TMS effects can be discussed it will also lead to more complex modelling of TMS  
2 effects on brain tissue.  
3  
4  
5  
6  
7

## 8 **References**

- 9  
10 Bestmann S, Ruff CC, Blankenburg F, Weiskopf N, Driver J, and Rothwell JC. Mapping causal  
11 interregional influences with concurrent TMS-fMRI. *Exp Brain Res* 191: 383-402, 2008.  
12  
13 Ermentrout GB, Galan RF, and Urban NN. Reliability, synchrony and noise. *Trends Neurosci* 31:  
14 428-434, 2008.  
15  
16 Green DM, Swets JA. *Signal Detection Theory and Psychophysics*. Wiley, New York 1966  
17  
18 Harris JA, Clifford CW, and Miniussi C. The functional effect of transcranial magnetic stimulation:  
19 signal suppression or neural noise generation? *J Cogn Neurosci* 20: 734-740, 2008a.  
20  
21 Harris IM, Benito CT, Ruzzoli M, and Miniussi C. Effects of Right Parietal Transcranial Magnetic  
22 Stimulation on Object Identification and Orientation Judgments. *J Cogn Neurosci* 20: 916-926,  
23 2008b.  
24  
25 Miniussi C, Cappa SF, Cohen L, Floel A, Fregni F, Nitsche M, Oliveri M, Pascual-Leone A, Paulus  
26 W, Priori A, Walsh V. Efficacy of rTMS/tDCS in cognitive neurorehabilitation. *Brain*  
27 Stimulation 1: 326-336, 2008.  
28  
29 Moss F, Ward LM, and Sannita WG. Stochastic resonance and sensory information processing: a  
30 tutorial and review of application. *Clin Neurophysiol* 115: 267-281, 2004.  
31  
32 Ruff CC, Driver J, and Bestmann S. Combining TMS and fMRI: From 'virtual lesions' to  
33 functional-network accounts of cognition. *Cortex*, in press.  
34  
35 Sack AT and Linden DE. Combining transcranial magnetic stimulation and functional imaging in  
36 cognitive brain research: possibilities and limitations. *Brain Res Rev* 43: 41-56, 2003.  
37  
38 Siebner HR, Hartwigsen G, Kassuba T, and Rothwell J. How does transcranial magnetic stimulation  
39 modify neuronal activity in the brain? Implications for studies of cognition. *Cortex*, in press.  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 Silvano J, Muggleton N, and Walsh V. State-dependency in brain stimulation studies of perception  
2 and cognition. Trends Cogn Sci 12: 447-454, 2008.  
3

4 Stein RB, Gossen ER, and Jones KE. Neuronal variability: noise or part of the signal? Nat Rev  
5 Neurosci 6: 389-397, 2005.  
6

7 Wagner T, Rushmore J, Eden U, and Valero-Cabre A. Biophysical foundations underlying TMS:  
8 Setting the stage for an effective use of neurostimulation in the cognitive neurosciences.  
9 Cortex, in press.  
10

11 Walsh V and Cowey A. Magnetic stimulation studies of visual cognition. Trends Cogn Sci 2: 103-  
12 110, 1998.  
13

14 Walsh V and Cowey A. Transcranial magnetic stimulation and cognitive neuroscience. Nat Rev  
15 Neurosci 1: 73-79, 2000.  
16

17 Walsh V and Pascual-Leone A. Transcranial Magnetic Stimulation: A Neurochronometrics of Mind.  
18 Cambridge, Massachusetts: Mit Press, 2003.  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65